

群感受野技术在感知觉的脑机制研究领

$$\begin{array}{c}
[s(x,y,t)] \\
\\
1 \\
0 \\
\text{pRF} \quad [g(x,y)] \\
[s(x,y,t)] \quad \text{pRF} \\
[r(t)] \\
r(t) = \underset{x,y}{s(x,y,t)} g(x,y) \\
(\text{hemo-}) \\
\text{dynamic response function, HRF) } h(t) \quad r(t) \\
\text{pRF} \quad \text{BOLD} \\
[p(t)] \\
p(t) = r(t) * h(t) \\
\\
\text{pRF} \quad \text{BOLD} \quad [p(t)] \\
\text{BOLD} \quad [y(t)] \quad \text{RSS} \\
\text{RSS} = \sum_i (y(t) - p(t)\beta)^2 \quad \text{RSS} \\
\text{pRF} \quad \text{pRF} \\
\text{pRF}
\end{array}$$

2 pRF技术的应用

fMRI pRF

$$\begin{array}{c}
 \text{pRF} \\
 |[1, 9, 14, 21, 22] \\
 \text{pRF} \\
 |[24] \\
 \text{pRF} \\
 |[6, 23, 28, 29] \\
 |[18, 19, 26] \\
 |[25]
 \end{array}$$

2.1 基于pRF位置信息的应用

2.4 pRF技术的泛化应用(非视觉的pRF技术、注意野技术和群调谐技术)



fMRI pRF

4 总结和展望

fMRI pRF

fMRI

pRF

pRF

pRF

pRF

pRF

pRF

pRF

Dumoulin Wandell

pRF

()

(

pRF

pRF

[1]

(7T)

fMRI

pRF

(

1 mm)

(

6

pRF

pRF

pRF

1

pRF

pRF

pRF

[g(x,y)]

Flom

1963

(visual crowding

(

effect

)

[38]

fMRI

pRF

pRF

pRF

pRF

pRF

pRF

pRF

pRF

pRF [9, 35, 36]

(model-free)

(

pRF

pRF^[37]

)

[39–44]

pRF

pRF

pRF


pRF

()

pRF

pRF

pRF



参考文献

- 1 Dumoulin SO, Wandell BA. Population receptive field estimates in human visual cortex. *Neuroimage* 2008; 39(2): 647–660.
- 2 Smith AT, Singh KD, Williams AL, Greenlee MW. Estimating receptive field size from fMRI data in human striate and extrastriate visual cortex. *Cereb Cortex* 2001; 11(12): 1182–1190.
- 3 Victor JD, Purpura K, Katz E, Mao B. Population encoding of spatial frequency, orientation, and color in macaque V1. *J Neurophysiol* 1994; 72(5): 2151–2166.
- 4 Wandell BA, Winawer J. Computational neuroimaging and population receptive fields. *Trends Cogn Sci* 2015; 19(6): 349–357.
- 5 Baseler HA, Gouws A, Haak KV, Racey C, Crossland MD, Tufail A, Rubin GS, Cornelissen FW, Morland AB. Large-scale remapping of visual cortex is absent in adult humans with macular degeneration. *Nat Neurosci* 2011; 14(5): 649–655.
- 6 Kok P, de Lange FP. Shape perception simultaneously up- and downregulates neural activity in the primary visual cortex. *Curr Biol* 2014; 24(13): 1531–1535.
- 7 Lee S, Papanikolaou A, Logothetis NK, Smirnakis SM, Kellis GA. A new method for estimating population receptive field topography C, i

- feld (pRF) measurements of chromatic responses in human visual cortex using fMRI. *Neuroimage* 2018; 167: 84–94.
- 27 Zhang X, Fang F. Cortical magnification factor and population receptive field size in human V1 predict the bottom-up saliency map. *J Vis* 2014; 14(10): 537. DOI: 10.1167/14.10.537.
- 28 Zhu Z, He D, Fang F. Crowding suppresses cortical responses to the target in human early visual cortex. *J Vis* 2015; 15(12): 101.
- 29 Zuiderbaan W, Harvey BM, Dumoulin SO. Image identification from brain activity using the population receptive field model. *PLoS One* 2017; 12(9): e0183295.
- 30 Deyoe EA, Carman GJ, Bandettini P, Glickman S, Wieser J, Cox R, Miller D, Neitz J. Mapping striate and extrastriate visual areas in human cerebral cortex. *Proc Natl Acad Sci U S A* 1996; 93(6): 2382–2386.
- 31 Dumoulin SO, Hoge RD, Baker CL Jr, Hess RF, Achtman RL, Evans AC. Automatic volumetric segmentation of human visual retinotopic cortex. *Neuroimage* 2003; 18(3): 576–587.
- 32 Engel SA, Glover GH, Wandell BA. Retinotopic organization in human visual cortex and the spatial precision of functional MRI. *Cereb Cortex* 1997; 7(2): 181–192.
- 33 Sereno MI, Dale AM, Reppas JB, Kwong KK, Belliveau JW, Brady TJ, Rosen BR, Tootell RB. Borders of multiple visual areas in humans revealed by functional magnetic resonance imaging. *Science* 1995; 268(5212): 889–893.
- 34 Sprague TC, Serences JT. Attention modulates spatial priority maps in the human occipital, parietal and frontal cortices. *Nat Neurosci* 2013; 16(12): 1879–1887.
- 35 Silson EH, Reynolds RC, Kravitz DJ, Baker CI. Differential sampling of visual space in ventral and dorsal early visual cortex. *J Neurosci* 2018; 38(9): 2294–2303.
- 36 Zeidman P, Silson EH, Schwarzkopf DS, Baker CI, Penny W. Bayesian population receptive field modelling. *Neuroimage* 2017; pii: S1053-8119(17)30746-2. doi: 10.1016/j.neuroimage.2017.09.008.
- 37 Greene CA, Dumoulin SO, Harvey BM, Ress D. Measurement of population receptive fields in human early visual cortex using back-projection tomography. *J Vis* 2014; 14(1): 1–7.
- 38 Flom MC, Heath GG, Takahashi E. Contour interaction and visual resolution: Contralateral effects. *Science* 1963; 142(3594): 979–980.
- 39 Anderson EJ, Dakin SC, Schwarzkopf DS, Rees G, Greenwood JA. The neural correlates of crowding-induced changes in appearance. *Curr Biol* 2012; 22(13): 1199–1206.
- 40 Bi T, Cai P, Zhou T, Fang F. The effect of crowding on orientation-selective adaptation in human early visual cortex. *J Vis* 2009; 9(11): 1–10.
- 41 Chen J, He Y, Zhu Z, Zhou T, Peng Y, Zhang X, Fang F. Attention-dependent early cortical suppression contributes to crowding. *J Neurosci* 2014; 34(32): 10465–10474.
- 42 Fang F, He S. Crowding alters the spatial distribution of attention modulation in human primary visual cortex. *J Vis* 2008; 8(9): 1–6.
- 43 Freeman J, Donner TH, Heeger DJ. Inter-area correlations in the ventral visual pathway reflect feature integration. *J Vis* 2011; 11(4): 74–76.
- 44 Millin R, Arman AC, Chung ST, Tjan BS. Visual crowding in V1. *Cereb Cortex* 2013; 24(12): 3107–3115.