

Enhancing visual perceptual learning using transcranial electrical stimulation: Transcranial alternating current stimulation outperforms both transcranial direct current and random noise stimulation

Qing He*



Xinyi Zhu*



Fang Fang



Diverse strategies can be employed to enhance visual skills, including visual perceptual learning (VPL) and transcranial electrical stimulation (tES). Combining VPL and tES is a popular method that holds promise for producing significant improvements in visual acuity within a short time frame. However, there is still a lack of comprehensive evaluation regarding the effects of combining different types of tES and VPL on enhancing visual function, especially with a larger sample size. In the present study, we recruited four groups of subjects

(26 subjects each) to learn an orientation discrimination task with five daily training sessions. During training, the occipital region of each subject was stimulated by one type of tES—anodal transcranial direct current stimulation (tDCS), alternating current stimulation (tACS) at 10 Hz, high-frequency random noise stimulation (trNS), and sham tACS—while the subject performed the training task. We found that, compared with the sham stimulation, both the high-frequency trNS and the 10-Hz tACS facilitated VPL efficiently in

Qing He, Xinyi Zhu, & Fang Fang (2023).

Journal of Vision, 23(14):2, 1–12, <https://doi.org/10.1167/jov.23.14.2>

Journal of

<https://doi.org/10.1167/jov.23.14.2>

Received August 28, 2023; published December 6, 2023

ISSN 1534-7362 Copyright 2023 The Authors



terms of learning rate and performance improvement, but there was little modulatory effect in the anodal tDCS condition. Remarkably, the 10-Hz tACS condition exhibited superior modulatory effects compared with the tRNS condition, demonstrating the strongest modulation among the most commonly used tES types for further enhancing vision when combined with VPL. Our results suggest that alpha oscillations play a vital role in VPL. Our study provides a practical guide for vision rehabilitation.

Introduction

Our visual system is remarkably malleable (i.e., has visual plasticity) throughout the entire life span, for both mature and degenerative brains. It has been well documented that various methods can bring about visual plasticity, such as extensive training and non-invasive brain stimulation techniques (He, Yang, Zhao, & Fang, 2022). Regarding visual plasticity induced by extensive training (i.e., visual perceptual learning [VPL]), many advancements have been made in unveiling its characteristics, brain loci, and neural manifestations over the last three decades (He, Yang, & Zhao, 2022). More importantly, VPL has been widely applied in various fields (Lu, Lin, & Doshier, 2016), such as low-vision rehabilitation (Huang et al., 2022; Levi & Polat, 1996), remediation for dyslexia (Gori & Facoetti, 2014; Meng, Lin, Wang, Jiang, & Song, 2014), x-ray security screening (McCarley, Kramer, Wickens, Vidoni, & Boot, 2004), sports and military training (Hadlow, Panchuk, Mann, Portus, & Abernethy, 2018), and medical imaging education (Alexander, Waite, Macknik, & Martinez-Conde, 2020). At the same time, our visual plasticity can also be induced by transcranial electrical stimulation (tES)—a non-invasive neuromodulatory technique in which weak electrical fields are delivered through electrodes positioned on the scalp surface. The three most common types of tES techniques are categorized based on the current waveform: transcranial direct current stimulation (tDCS) (Nitsche & Paulus, 2000; Priori, Berardelli, Rona, Accornero, & Manfredi, 1998), transcranial alternating current stimulation (tACS) (Antal et al., 2008), and transcranial random noise stimulation (tRNS) (Terney, Chaieb, Moliadze, Antal, & Paulus, 2008). All of these three types of tES techniques have been found to be able to modify cortical excitability and modulate visual functions in both healthy and clinical populations (Bello et al., 2023), such as contrast sensitivity (Antal, Nitsche, & Paulus, 2001; Potok et al., 2023), visual acuity (Bocci et al., 2018; Reinhart, Xiao, McClenahan, & Woodman, 2016), motion direction discrimination (Battaglini et al., 2023; Ghin, Pavan, Contillo, & Mather, 2018), and peripheral target

identification in a crowded environment (Battaglini, Ghiani, Casco, & Ronconi, 2020; Chen, Zhu, He, & Fang, 2021), thus establishing causal links between brain activity and cognitive functions (Zhang, Zhang, Cai, Luo, & Fang, 2019).

Currently, there is an increasing focus on modulating VPL by tES, which is both theoretically significant and practically useful. Theoretically, through selective modulation of the brain during distinct phases of learning, mechanisms that support successful task acquisition and consolidation may be more fully characterized (He, Yang, & Fang, 2021; He, Yang, Gong, Bi, & Fang, 2022; He, Yang, & Zhao, 2022; Wu et al., 2023; Yang, He, & Fang, 2022). From the perspective of translational applications, attaining the greatest training benefits effect in the briefest time frame provides most benefit to subjects, whereas in a typical practical-oriented application case thousands of trials across multiple training sessions are usually required. Therefore, boosting VPL by tES is expected to accelerate learning and allow subjects to obtain greater benefits (He, Yang, & Gong, 2022; Herpich et al., 2019).

Different types of tES have been adopted to modulate VPL (for a review, see He, Yang, & Zhao, 2022). With regard to the effect of tDCS on modulating VPL, the results are mixed, and no consistent result has been found regardless of whether tDCS was administered before task execution (offline mode) or during task execution (online mode). For example, when tDCS was applied during task execution, anodal tDCS was found to be effective in boosting VPL in some studies (Frangou, Correia, & Kourtzi, 2018; Wu et al., 2023), but in other studies no obvious modulatory effect was found (Fertonani, Pirulli, & Miniussi, 2011; Herpich et al., 2019; Larcombe, Kennard, O'Shea, & Bridge, 2018; Larcombe et al., 2018; Wu et al., 2022), and even a suppressive effect was observed (Jia et al., 2022b). Studies using offline tDCS in VPL tasks showed similarly conflicting results (Pirulli, Fertonani, & Miniussi, 2013; Pirulli, Fertonani, & Miniussi, 2014; Wu et al., 2023). By contrast, the results of modulating VPL by tRNS are relatively consistent. Specifically, high-frequency (100–640 Hz) tRNS boosts VPL effectively in both healthy and clinical populations (Camilleri, Pavan, Ghin, Battaglini, & Campana, 2014; Campana, Camilleri, Pavan, Veronese, & Lo Giudice, 2014; Cappelletti, Pikkat, Upstill, Speekenbrink, & Walsh, 2015; Conto et al., 2021; Donkor et al., 2021). With regard to tACS, although the underlying neural mechanisms of action of tACS on the brain are relatively clear among tES techniques (Johnson et al., 2020; Krause, Vieira, Csorba, Pilly, & Pack, 2019; Zaehle, Rach, & Herrmann, 2010), only limited studies have been conducted to modulate VPL. Specifically, when we previously stimulated subjects' visual cortical areas with tACS at different frequencies, we found that only occipital 10-Hz tACS was able to boost VPL,

and no such effect was found for other stimulation frequencies (e.g., 6, 20, and 40 Hz) (He, Yang, & Gong, 2022). Similarly, in another study, occipital 3-Hz tACS was found to show little effect on VPL (Zizlsperger, Kummel, & Haarmeier, 2016). Moreover, we also found that the modulatory effect of 10-Hz tACS was absent when other cortical areas were stimulated (He, Yang, & Gong, 2022). Altogether, these studies demonstrate that tACS facilitates VPL in a frequency- and location-specific manner.

Despite advancements in VPL being observed following the application of diverse types of tES, a systematic evaluation of the modulatory effects of tES on VPL is still lacking. Notably, studies on modulating VPL by tES usually employ a relatively small sample size design, raising the risk of sampling bias and limiting the generalizability of those findings (Minarik et al., 2016). We recruited four groups of subjects to learn an orientation discrimination task, and each group received a specific form of stimulation, including sham, tDCS, high-frequency tRNS, and 10-Hz tACS during their training phase. Notably, there were 26 participants in each group, which provided sufficient statistical power and representativeness to ensure the reliability and generalizability of our findings.

Methods


Subjects

A total of 104 right-handed healthy adults with normal or correct-to-normal vision took part in the present study. Subjects were assigned to one of four conditions (26 subjects in each condition): anodal tDCS (11 females; mean age, 21.54 ± 2.42 years), high-frequency tRNS (20 females; mean age, 21.77 ± 2.21 years), 10-Hz tACS (18 females; mean age, 21.15 ± 4.58 years), and sham 10-Hz tACS (17 females; mean age, 22.04 ± 2.89 years). The sample sizes were determined based on our previous study (alpha level = 0.05, power = 80%, two-tailed Cohen's $d = 0.96$) (He, Yang, & Gong, 2022). Specifically, all data from that study (He, Yang, & Gong, 2022) for the 10-Hz tACS condition ($n = 21$) and the sham 10-Hz tACS condition ($n = 20$) were included in the present study, and all other data were newly collected. We recruited five more subjects for the 10-Hz tACS condition and six more subjects for the sham stimulation condition in the present study to obtain more robust results. A screening questionnaire was administered for each subject before starting the study. Subjects were excluded if they met the following criteria: (1) age older than 30 years or younger than 18 years, (2) a history of neural surgery or epileptic seizures or any psychiatric or neurological disorders, (3) sleep disorders or a

total sleep time less than 7 hours per night over the last 2 weeks, or (4) during the ovulation phase of the menstrual cycle or pregnancy (He et al., 2019; He, Yang, & Gong, 2022). All experimental protocols and procedures were approved by the Ethics Committee of the School of Psychological and Cognitive Sciences at Peking University. Before participation, informed written consent was obtained from each subject.

Apparatus and stimulation protocol

MATLAB R2015a (MathWorks, Natick, MA) and Psychtoolbox-3 extensions (Brainard, 1997) were used to generate and control visual stimuli, which were presented on a liquid crystal display (LCD) monitor (Display ++ LCD Monitor; Cambridge Research Systems, Rochester, UK) with a gray background (mean luminance = 30 cd/m^2 , width = 70 cm, spatial resolution = 1920×1080 pixels, refresh rate = 120 Hz). The only source of light in the room was the monitor. The head of each subject was fixed on a chin and head rest at a viewing distance of 70 cm. To ensure that the subject's eye position was stable within 1° from the fixation point when visual stimuli were presented, the EyeLink 1000 Plus eye-tracking system (SR Research Ltd., Ottawa, ON, Canada) was used to monitor subjects' eye movements throughout the entire experiment.



of the pixels in these patches were replaced with random noise) were presented in the lower left quadrant of the visual field, 5° from the fixation point. In each trial, a small fixation point was presented first for 500 ms, followed by two Gabor patches with orientations of 26° and $26^\circ + \theta$, which appeared 100 ms each in a random order with a 500-ms blank interval (Figure 1A). A two-alternative forced-choice (2AFC) method was used in the task, and subjects were instructed to judge the orientation change of the second Gabor patch relative to the first one (counterclockwise or clockwise) by pressing keys. Subjects' orientation discrimination thresholds at 75% accuracy were estimated using a QUEST staircase procedure, such that the θ varied trial by trial (Watson & Pelli, 1983). Subjects rested after each staircase. No feedback was provided in all test and training sessions.

Design

A single-blind, sham-controlled, between-subjects design was adopted to explore the modulatory effect of different types of tES on orientation discrimination learning. Subjects were trained on the orientation discrimination task for five consecutive days. Before and after the five daily training sessions, test sessions were conducted: pretraining (Pre) and post-training (Post) (Figure 1B). Each test session and each training session consisted of six and nine QUEST staircases of 50 trials, respectively.

The hemisphere contralateral to the visual field where the visual stimuli were presented was stimulated. Two electrodes were placed over each subject's visual cortex and vertex (i.e., O2 and Cz in the international 10–20 EEG system, respectively) (Figure 1C). Electrical

stimulation was delivered concurrently with training, with a ramp-up of 30 seconds at the beginning of each training session.

Statistical analysis

The threshold for each session was estimated by calculating the geometric mean of thresholds from all QUEST staircases in that session. The performance change after training was quantified by calculating the percent improvement as $[(\text{pretraining threshold} - \text{post-training threshold})/\text{pretraining threshold}] \times 100\%$. All estimated thresholds were then normalized: The estimated threshold for each session was divided by the estimated threshold at Pre and then multiplied by 100%. In order to describe the process of threshold change during the learning course, we fitted the learning curves of normalized orientation discrimination thresholds across all sessions using a power function:

$$\log_{10}(Th(t)) = \rho \times \log_{10}(t)$$

In this function, Th represents the predicted normalized threshold, t is the number of training sessions, and ρ is the learning rate (Yang et al., 2020). The threshold declines with training, such that the value of ρ should be negative, and a smaller value indicates a faster learning speed. To minimize the sum of squared differences between model predictions and observed values, a nonlinear, least-square method was implemented in MATLAB.

A mixed-design analysis of variance (ANOVA) using *Condition* as a between-subjects factor (tDCS, tRNS, tACS, and sham) was used to analyze the orientation discrimination thresholds. Raw thresholds at Pre, learning rates, and percent improvements were analyzed by ANOVA using *Condition* as a between-subjects factor. We used the Benjamini–Hochberg method to control the false discovery rate for multiple comparisons. Partial eta squared (η_p^2) and Cohen's d were calculated to measure the effect size for ANOVAs and t -tests. Statistical analyses were conducted using R (R Foundation for Statistical Computing, Vienna, Austria) (R Core Team, 2023).

Results

Training improved task performance substantially

First, a one-way ANOVA with *Condition* (tDCS, tRNS, tACS, and sham) as a between-subjects factor

on the thresholds at Pre was conducted to examine the baseline variance among all enrolled conditions. The statistical results showed that the main effect of *Condition* was not significant, $F(3, 100) = 0.37, p = 0.79, \eta_p^2 = 0.01$, demonstrating that subjects had comparable baseline performance across all stimulation conditions (Figure 2A). Next, to assess the learning effect on the trained task, we employed a mixed-design ANOVA that incorporated *Session* (Pre and Post) as a within-subjects factor, *Condition* as a between-subjects factor, and the subjects' orientation discrimination thresholds as the dependent variable. The statistical results showed that the main effect of *Session*, $F(1, 100) = 207.89, p < 0.001, \eta_p^2 = 0.68$, and the interaction between *Session* and *Condition*, $F(3, 100) = 3.29, p = 0.02, \eta_p^2 = 0.09$, were significant, but the main effect of *Condition*, $F(3, 100) = 0.95, p = 0.42$, was not significant, demonstrating that the subjects' performance improved after training and the performance improvement was different across all stimulation conditions. Further analyses showed that the thresholds at Post were lower than those at Pre for all stimulation conditions: for sham, $t(25) = 4.33, p_{\text{adj}} < 0.001$, Cohen's $d = 0.85$; for tDCS, $t(25) = 6.77, p_{\text{adj}} < 0.001$, Cohen's $d = 1.33$; for tRNS, $t(25) = 10.21, p_{\text{adj}} < 0.001$, Cohen's $d = 2.00$; for tACS, $t(25) = 9.01, p_{\text{adj}} < 0.001$, Cohen's $d = 1.77$ (Figure 2B). Importantly, no feedback was provided during the training course, suggesting that subjects acquired the improved ability to discriminate the trained stimuli in an unsupervised learning manner (Frank et al., 2020; Tsodyks & Gilbert, 2004).

Learning rate was modulated by tES

The efficiency of performance improvement in learning is typically quantified by learning rate, which in this case is represented by the ρ value. We examined ρ -value differences among the different stimulation conditions. A one-way ANOVA revealed that the main effect of *Condition* on ρ value was significant, $F(3, 100) = 9.20, p < 0.001, \eta_p^2 = 0.22$, demonstrating that subjects in different stimulation conditions showed different learning rates (Figure 2C). Specifically, pairwise t -tests showed that there was no significant difference between the sham condition and the tDCS condition in learning rate, $t(50) = 0.90, p_{\text{adj}} = 0.38$, Cohen's $d = 0.25$, whereas the learning rates in both the tRNS and the tACS conditions were significantly greater than that in the sham condition: for tRNS versus sham, $t(50) = 2.90, p_{\text{adj}} = 0.01$, Cohen's $d = 0.80$; for tACS versus sham, $t(50) = 4.44, p_{\text{adj}} < 0.001$, Cohen's $d = 1.23$. Compared with the tDCS condition, the learning rate in the tACS condition was higher, $t(50) = 3.76, p_{\text{adj}} = 0.001$, Cohen's $d = 1.12$. The learning rate in the tRNS condition was also higher than that in the tDCS condition, $t(50) = 19.2, p_{\text{adj}} < 0.001$, Cohen's $d = 5.72$. The learning rate in the tACS condition was also higher than that in the tRNS condition, $t(50) = 19.2, p_{\text{adj}} < 0.001$, Cohen's $d = 5.72$.

discrimination learning, in terms of both learning rate and overall performance improvement achieved. Finally, occipital 10-Hz tACS showed the best modulatory effects.

Discussion

In the present study, we evaluated the effect of several of the most common types of tES techniques (i.e., tDCS, tRNS, and 10-Hz tACS) on the further enhancement of vision by modulating VPL. Our results revealed distinct modulatory effects of tES with different current forms when applied during learning on the orientation discrimination task. Specifically, compared with the sham stimulation, anodal tDCS showed little effect on modulating VPL, but both high-frequency tRNS and 10-Hz tACS were effective in boosting VPL. Additionally, 10-Hz tACS exhibited a superior modulatory effect compared to the other current stimulation forms. To the best of our knowledge, the current study is the first to systematically evaluate the effects of various types of tES techniques on modulating VPL. Importantly, compared with previous studies investigating the modulatory effects of tES on VPL, the present study has the largest sample size (~30 subjects per condition), thus ensuring sufficient statistical power for between-condition comparisons. In short, our findings will provide guidance for translational applications of combining VPL and tES, as well as insights into the neural mechanisms of VPL.

We found that anodal tDCS applied concurrently with training had no effect on modulating VPL, which is consistent with previous studies in short-term orientation discrimination learning (Fertonani et al., 2011) or in multi-session learning on motion direction discrimination (Fertonani, Pirulli, Bollini, Miniussi, & Bortoletto, 2019; Herpich et al., 2019; Larcombe & Kennard, 2018; Larcombe & Kulyomina, 2018; Wu et al., 2022). However, our results are not consistent with some studies in which anodal tDCS was effective in facilitating VPL (Olma et al., 2013; Sczesny-Kaiser et al., 2016; Van Meel, Daniels, de Beeck, & Baeck, 2016; Wu et al., 2023), or even impairing VPL with a short training session (Grasso, Tonolli, Bortoletto, & Miniussi, 2021; Grasso, Tonolli, & Miniussi, 2020; Jia et al., 2022a; Learmonth, Thut, Benwell, & Harvey, 2015). The discrepancies among these studies mentioned above can be caused by many factors, such as training tasks, training regimes, and stimulation settings. Of note, the small sample sizes adopted in previous studies may have biased the results (Minarik et al., 2016). Here, using a relatively large sample size design, we found that applying anodal tDCS concurrently during training had a negligible effect on enhancing VPL. The null effect

of anodal tDCS on modulating VPL during training might be caused by the timing of the stimulation applied. Previous studies have found that anodal tDCS applied prior to task execution (i.e., the offline mode) was effective in boosting VPL (Pirulli et al., 2013), but no such modulatory effect was found when anodal tDCS was administered during performance of the orientation discrimination task (i.e., the online mode) (Pirulli et al., 2013), just like the findings in the present study.

We found that high-frequency tRNS was effective in boosting VPL, which is consistent with previous studies (Camilleri et al., 2014; Camilleri, Pavan, & Campana, 2016; Contemori, Trotter, Cottreau, & Maniglia, 2019;

spike-timing (Huang et al., 2021; Johnson et al., 2020; Krause et al., 2019), such that attention-related cognitive processing is strengthened by occipital tACS at 10 Hz directly (Clayton, Yeung, & Cohen Kadosh, 2019). In contrast, high-frequency tRNS over the occipital regions may not modulate attentional processing efficiently. Previous studies have shown that applying high-frequency tRNS over the parietal regions can improve visual attention (Edwards, Contò, Bucci, & Battelli, 2020; Shalev, De Wandel, Dockree, Demeyere, & Chechlacz, 2018; Tyler, Conto, & Battelli, 2018), but no such effect was observed when stimulating the visual areas (Conto et al., 2021; Conto, Tyler, Paletta, & Battelli, 2023; Tyler et al., 2018). Therefore, the superior modulatory effects of occipital 10-Hz tACS on VPL compared to high-frequency tRNS may be attributed to different efficacies to mediate attentional processes. Here, this explanation is referred to as the efficacy of attention hypothesis, which requires further investigation in the future.

Our findings will provide a practical guide for vision rehabilitation. VPL has been widely adopted to restore low-vision in neuroophthalmology (Lu et al., 2016). In practice, it is a common desire of patients and their families to achieve the best effect in a short course of treatment (He, Yang, & Gong, 2022). However, typical clinic-oriented studies require weeks to months of training and consist of thousands of trials, placing a significant economic and psychological burden on both the patients' families and society in general (Herpich et al., 2019). Therefore, developing methods that can speed up the process and improve the treatment effectiveness is currently a hot research topic. Our results show that 10-Hz tACS over visual areas during training accelerated the learning process and maximized learning gains in healthy adults, findings that are expected to be significant for patients who need to learn or relearn skills due to injuries or illnesses. Overall, our findings have the potential to significantly impact the field of vision habilitation (Herpich et al., 2019; Huang et al., 2022; Wu et al., 2023), and further studies will help to fully determine the effectiveness of this method for patients. For example, researchers have found that patients with cortical blindness (vision loss caused by damage to the primary visual cortex) regained the ability to discriminate visual motion direction after extensive training concurrent with high-frequency tRNS (Herpich et al., 2019). The current study, however, has potential limitations with regard to clinical applications. For example, in this study, the majority of the participants were under the age of 30 years, and it is unclear if the protocols employed would be applicable to other populations (e.g., those of an elderly age) or other training tasks.

Keywords: transcranial electrical stimulation, tDCS, tACS, tRNS, visual perceptual learning, visual plasticity, unsupervised learning, implicit learning

Acknowledgments

Supported by the National Science and Technology Innovation 2030 Major Program (2022ZD0204802) and the National Natural Science Foundation of China (31930053). QH was supported in part by a Postdoctoral Fellowship of the Peking-Tsinghua Center for Life Sciences.

Commercial relationships: none.

Corresponding author: Fang Fang.

Email: ffang@pku.edu.cn.

Address: School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China.

*QH and XZ contributed equally to this article.

References

- Alexander, R. G., Waite, S., Macknik, S. L., & Martinez-Conde, S. (2020). What do radiologists look for? Advances and limitations of perceptual learning in radiologic search. *Journal of Vision*, 20(10):17, 1–13, <https://doi.org/10.1167/jov.20.10.17>.
- Antal, A., Boros, K., Poreisz, C., Chaieb, L., Terney, D., & Paulus, W. (2008). Comparatively weak after-effects of transcranial alternating current stimulation (tACS) on cortical excitability in humans. *Brain Stimulation*, 1(2), 97–105, <https://doi.org/10.1016/j.brs.2007.10.001>.
- Antal, A., Nitsche, M. A., & Paulus, W. (2001). External modulation of visual perception in humans. *NeuroReport*, 12(16), 3553–3555, <https://doi.org/10.1097/00001756-200111160-00036>.
- Battaglini, L., Casco, C., Fertoni, A., Miniussi, C., Di Ponzio, M., & Vicovaro, M. (2023). Noise in the brain: Transcranial random noise stimulation and perceptual noise act on a stochastic resonance-like mechanism. *European Journal of Neuroscience*, 57(12), 2097–2111, <https://doi.org/10.1111/ejn.15965>.
- Battaglini, L., Ghiani, A., Casco, C., & Ronconi, L. (2020). Parietal tACS at beta frequency improves vision in a crowding regime. *NeuroImage*, 208, 116451, <https://doi.org/10.1016/j.neuroimage.2019.116451>.
- Bays, B. C., Visscher, K. M., Le Dantec, C. C., & Seitz, A. R. (2015). Alpha-band EEG activity in perceptual learning. *Journal of Vision*, 15(10):7, 1–12, <https://doi.org/10.1167/15.10.7>.
- Bello, U. M., Wang, J., Park, A. S. Y., Tan, K. W. S., Cheung, B. W. S., Thompson, B., . . .

- Cheong, A. M. Y. (2023). Can visual cortex non-invasive brain stimulation improve normal visual function? A systematic review and meta-analysis. *Frontiers in Neuroscience*, *17*, 1119200, <https://doi.org/10.3389/fnins.2023.1119200>.
- Bocci, T., Nasini, F., Caleo, M., Restani, L., Barloscio, D., Ardolino, G., . . . Sartucci, F. (2018). Unilateral application of cathodal tDCS reduces transcallosal inhibition and improves visual acuity in amblyopic patients. *Frontiers in Behavioral Neuroscience*, *12*, 109, <https://doi.org/10.3389/fnbeh.2018.00109>.
- Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436, <https://doi.org/10.1163/156856897X00357>.
- Camilleri, R., Pavan, A., & Campana, G. (2016). The application of online transcranial random noise stimulation and perceptual learning in the improvement of visual functions in mild myopia. *Neuropsychologia*, *89*, 225–231, <https://doi.org/10.1016/j.neuropsychologia.2016.06.024>.
- Camilleri, R., Pavan, A., Ghin, F., Battaglini, L., & Campana, G. (2014). Improvement of uncorrected visual acuity and contrast sensitivity with perceptual learning and transcranial random noise stimulation in individuals with mild myopia. *Frontiers in Psychology*, *5*, 1234, <https://doi.org/10.3389/fpsyg.2014.01234>.
- Campana, G., Camilleri, R., Pavan, A., Veronese, A., & Lo Giudice, G. (2014). Improving visual functions in adult amblyopia with combined perceptual training and transcranial random noise stimulation (tRNS): A pilot study. *Frontiers in Psychology*, *5*, 1402, <https://doi.org/10.3389/fpsyg.2014.01402>.
- Cappelletti, M., Pikkat, H., Upstill, E., Speekenbrink, M., & Walsh, V. (2015). Learning to integrate versus inhibiting information is modulated by age. *Journal of Neuroscience*, *35*(5), 2213–2225, <https://doi.org/10.1523/JNEUROSCI.1018-14.2015>.
- Chen, G., Zhu, Z., He, Q., & Fang, F. (2021). Offline transcranial direct current stimulation improves the ability to perceive crowded targets. *Journal of Vision*, *21*(2), 1–10, <https://doi.org/10.1167/jov.21.2.1>.
- Clayton, M. S., Yeung, N., & Cohen Kadosh, R. (2019). Electrical stimulation of alpha oscillations stabilizes performance on visual attention tasks. *Journal of Experimental Psychology: General*, *148*(2), 203–220, <https://doi.org/10.1037/xge0000502>.
- Contemori, G., Trotter, Y., Cottureau, B. R., & Maniglia, M. (2019). tRNS boosts perceptual learning in peripheral vision. *Neuropsychologia*, *125*, 129–136, <https://doi.org/10.1016/j.neuropsychologia.2019.02.001>.
- Conto, F., Edwards, G., Tyler, S., Parrott, D., Grossman, E., & Battelli, L. (2021). Attention network modulation via tRNS correlates with attention gain. *eLife*, *10*, e63782, <https://doi.org/10.7554/eLife.63782>.
- Conto, F., Tyler, S., Paletta, P., & Battelli, L. (2023). The role of the parietal lobe in task-irrelevant suppression during learning. *Brain Stimulation*, *16*(3), 715–723, <https://doi.org/10.1016/j.brs.2023.04.006>.
- Donkor, R., Silva, A. E., Teske, C., Wallis-Duffy, M., Johnson, A. P., & Thompson, B. (2021). Repetitive visual cortex transcranial random noise stimulation in adults with amblyopia. *Scientific Reports*, *11*(1), 3029, <https://doi.org/10.1038/s41598-020-80843-8>.
- Edwards, G., Contò, F., Bucci, L. K., & Battelli, L. (2020). Controlling brain state prior to stimulation of parietal cortex prevents deterioration of sustained attention. *Cerebral Cortex Communications*, *1*(1), tgaa069, <https://doi.org/10.1093/texcom/tgaa069>.
- Fertonani, A., Pirulli, C., Bollini, A., Miniussi, C., & Bortoletto, M. (2019). Age-related changes in cortical connectivity influence the neuromodulatory effects of transcranial electrical stimulation. *Neurobiology of Aging*, *82*, 77–87, <https://doi.org/10.1016/j.neurobiolaging.2019.07.009>.
- Fertonani, A., Pirulli, C., & Miniussi, C. (2011). Random noise stimulation improves neuroplasticity in perceptual learning. *Journal of Neuroscience*, *31*(43), 15416–15423, <https://doi.org/10.1523/JNEUROSCI.2002-11.2011>.
- Frangou, P., Correia, M., & Kourtzi, Z. (2018). GABA, not BOLD, reveals dissociable learning-dependent plasticity mechanisms in the human brain. *eLife*, *7*, e35854, <https://doi.org/10.7554/eLife.35854>.
- Frank, S. M., Qi, A., Ravasio, D., Sasaki, Y., Rosen, E. L., & Watanabe, T. (2020). Supervised learning occurs in visual perceptual learning of complex natural images. *Current Biology*, *30*(15), 2995–3000.e3, <https://doi.org/10.1016/j.cub.2020.05.050>.
- Ghin, F., Pavan, A., Contillo, A., & Mather, G. (2018). The effects of high-frequency transcranial random noise stimulation (hf-tRNS) on global motion processing: An equivalent noise approach. *Brain Stimulation*, *11*(6), 1263–1275, <https://doi.org/10.1016/j.brs.2018.07.048>.
- Gori, S., & Facoetti, A. (2014). Perceptual learning as a possible new approach for remediation and prevention of developmental dyslexia. *Vision Research*, *99*, 78–87, <https://doi.org/10.1016/j.visres.2013.11.011>.
- Grasso, P. A., Tonolli, E., Bortoletto, M., & Miniussi, C. (2021). tDCS over posterior parietal cortex increases cortical excitability but decreases learning: An ERPs

- and TMS-EEG study. *Brain Research*, 1753, 147227, <https://doi.org/10.1016/j.brainres.2020.147227>.
- Grasso, P. A., Tonolli, E., & Miniussi, C. (2020). Effects of different transcranial direct current stimulation protocols on visuo-spatial contextual learning formation: Evidence of homeostatic regulatory mechanisms. *Scientific Reports*, 10(1), 4622, <https://doi.org/10.1038/s41598-020-61626-7>.
- Hadlow, S. M., Panchuk, D., Mann, D. L., Portus, M. R., & Abernethy, B. (2018). Modified perceptual training in sport: A new classification framework. *Journal of Science and Medicine in Sport*, 21(9), 950–958, <https://doi.org/10.1016/j.jsams.2018.01.011>.
- He, Q., Lin, B. R., Zhao, J., Shi, Y. Z., Yan, F. F., & Huang, C. B. (2019). No effects of anodal transcranial direct current stimulation on contrast sensitivity function. *Restorative Neurology and Neuroscience*, 37(2), 109–118, <https://doi.org/10.3233/RNN-180881>.
- He, Q., Yang, X.-Y., & Fang, F. (2021). The role of anodal transcranial direct current stimulation (tDCS) in the consolidation of visual perceptual learning is mediated by the wake/sleep cycle. *Journal of Vision*, 21(9), 2346–2346, <https://doi.org/10.1167/jov.21.9.2346>.
- He, Q., Yang, X. Y., Gong, B., Bi, K., & Fang, F. (2022). Boosting visual perceptual learning by transcranial alternating current stimulation over the visual cortex at alpha frequency. *Brain Stimulation*, 15(3), 546–553, <https://doi.org/10.1016/j.brs.2022.02.018>.
- He, Q., Yang, X.-Y., Zhao, D., & Fang, F. (2022). Enhancement of visual perception by combining transcranial electrical stimulation and visual perceptual training. *Medical Review*, 2(3), 271–284, <https://doi.org/10.1515/mr-2022-0010>.
- Herpich, F., Melnick, M. D., Agosta, S., Huxlin, K. R., Tadin, D., & Battelli, L. (2019). Boosting learning efficacy with noninvasive brain stimulation in intact and brain-damaged humans. *Journal of Neuroscience*, 39(28), 5551–5561, <https://doi.org/10.1523/JNEUROSCI.3248-18.2019>.
- Huang, W. A., Stitt, I. M., Negahbani, E., Passey, D. J., Ahn, S., Davey, M., . . . Frohlich, F. (2021). Transcranial alternating current stimulation entrains alpha oscillations by preferential phase synchronization of fast-spiking cortical neurons to stimulation waveform. *Nature Communications*, 12(1), 3151, <https://doi.org/10.1038/s41467-021-23021-2>.
- Huang, X., Xia, H., Zhang, Q., Blakemore, C., Nan, Y., Wang, W., . . . Pu, M. (2022). New treatment for amblyopia based on rules of synaptic plasticity: A randomized clinical trial. *Science China Life Sciences*, 65(3), 451–465, <https://doi.org/10.1007/s11427-021-2030-6>.
- Jia, K., Frangou, P., Karlaftis, V. M., Ziminski, J. J., Giorgio, J., Rideaux, R., . . . Kourtzi, Z. (2022a). Neurochemical and functional interactions for improved perceptual decisions through training. *Journal of Neurophysiology*, 127(4), 900–912, <https://doi.org/10.1152/jn.00308.2021>.
- Jia, K., Frangou, P., Karlaftis, V. M., Ziminski, J. J., Giorgio, J., Rideaux, R., . . . Kourtzi, Z. (2022b). Neurochemical and functional interactions for improved perceptual decisions through training. *Journal of Neurophysiology*, 127(4), 900–912, <https://doi.org/10.1152/jn.00308.2021>.
- Johnson, L., Alekseichuk, I., Krieg, J., Doyle, A., Yu, Y., Vitek, J., . . . Opitz, A. (2020). Dose-dependent effects of transcranial alternating current stimulation on spike timing in awake nonhuman primates. *Science Advances*, 6(36), eaaz2747, <https://doi.org/10.1126/sciadv.aaz2747>.
- Krause, M. R., Vieira, P. G., Csorba, B. A., Pilly, P. K., & Pack, C. C. (2019). Transcranial alternating current stimulation entrains single-neuron activity in the primate brain. *Proceedings of the National Academy of Sciences, USA*, 116(12), 5747–5755, <https://doi.org/10.1073/pnas.1815958116>.
- Larcombe, S. J., Kennard, C., O’Shea, J., & Bridge, H. (2018). No effect of anodal transcranial direct current stimulation (tDCS) over hMT+ on motion perception learning. *Frontiers in Neuroscience*, 12, 1044, <https://doi.org/10.3389/fnins.2018.01044>.
- Larcombe, S. J., Kulyomina, Y., Antonova, N., Ajina, S., Stagg, C. J., Clatworthy, P. L., . . . Bridge, H. (2018). Visual training in hemianopia alters neural activity in the absence of behavioural improvement: A pilot study. *Ophthalmic and Physiological Optics*, 38(5), 538–549, <https://doi.org/10.1111/opo.12584>.
- Learmonth, G., Thut, G., Benwell, C. S., & Harvey, M. (2015). The implications of state-dependent tDCS effects in aging: Behavioural response is determined by baseline performance. *Neuropsychologia*, 74, 108–119, <https://doi.org/10.1016/j.neuropsychologia.2015.01.037>.
- Levi, D. M., & Polat, U. (1996). Neural plasticity in adults with amblyopia. *Proceedings of the National Academy of Sciences, USA*, 93(13), 6830–6834, <https://doi.org/10.1073/pnas.93.13.6830>.
- Lu, Z. L., Lin, Z., & Doshier, B. A. (2016). Translating perceptual learning from the laboratory to applications. *Trends in Cognitive Sciences*, 20(8), 561–563, <https://doi.org/10.1016/j.tics.2016.05.007>.
- McCarley, J. S., Kramer, A. F., Wickens, C. D., Vidoni, E. D., & Boot, W. R. (2004). Visual skills in airport-security screening. *Psychological Science*,

- 15(5), 302–306, <https://doi.org/10.1111/j.0956-7976.2004.00673.x>.
- Meng, X., Lin, O., Wang, F., Jiang, Y., & Song, Y. (2014). Reading performance is enhanced by visual texture discrimination training in Chinese-speaking children with developmental dyslexia. *PLoS One*, *9*(9), e108274, <https://doi.org/10.1371/journal.pone.0108274>.
- Michael, E., Covarrubias, L. S., Leong, V., & Kourtzi, Z. (2023). Learning at your brain's rhythm: Individualized entrainment boosts learning for perceptual decisions. *Cerebral Cortex*, *33*(9), 5382–5394, <https://doi.org/10.1093/cercor/bhac426>.
- Minarik, T., Berger, B., Althaus, L., Bader, V., Biebl, B., Brotzeller, F., . . . Sauseng, P. (2016). The importance of sample size for reproducibility of tDCS effects. *Frontiers in Human Neuroscience*, *10*, 453, <https://doi.org/10.3389/fnhum.2016.00453>.
- Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *Journal of Physiology*, *527*(Pt 3), 633–639, <https://doi.org/10.1111/j.1469-7793.2000.t01-1-00633.x>.
- Olma, M. C., Dargie, R. A., Behrens, J. R., Kraft, A., Irlbacher, K., Fahle, M., . . . Brandt, S. A. (2013). Long-term effects of serial anodal tDCS on motion perception in subjects with occipital stroke measured in the unaffected visual hemifield. *Frontiers in Human Neuroscience*, *7*, 314, <https://doi.org/10.3389/fnhum.2013.00314>.
- Pirulli, C., Fertonani, A., & Miniussi, C. (2013). The role of timing in the induction of neuromodulation in perceptual learning by transcranial electric stimulation. *Brain Stimulation*, *6*(4), 683–689, <https://doi.org/10.1016/j.brs.2012.12.005>.
- Pirulli, C., Fertonani, A., & Miniussi, C. (2014). Is neural hyperpolarization by cathodal stimulation always detrimental at the behavioral level? *Frontiers in Behavioral Neuroscience*, *8*, 226, <https://doi.org/10.3389/fnbeh.2014.00226>.
- Potok, W., Post, A., Beliaeva, V., Bachinger, M., Cassara, A. M., Neufeld, E., . . . Wenderoth, N. (2023). Modulation of visual contrast sensitivity with tRNS across the visual system, evidence from stimulation and simulation. *eNeuro*, *10*(6), ENEURO.0177–0122.2023, <https://doi.org/10.1523/ENEURO.0177-22.2023>.
- Priori, A., Berardelli, A., Rona, S., Accornero, N., & Manfredi, M. (1998). Polarization of the human motor cortex through the scalp. *NeuroReport*, *9*(10), 2257–2260, <https://doi.org/10.1097/00001756-199807130-00020>.
- R Core Team. (2023). *R: A Language and Environment for Statistical Computing (Version 4.2.3)*. Vienna, Austria: R Foundation for Statistical Computing.
- Reinhart, R. M. G., Xiao, W. X., McClenahan, L. J., & Woodman, G. F. (2016). Electrical stimulation of visual cortex can immediately improve spatial vision. *Current Biology*, *26*(14), 1867–1872, <https://doi.org/10.1016/j.cub.2016.05.019>.
- Szczesny-Kaiser, M., Beckhaus, K., Dinse, H. R., Schwenkreis, P., Tegenthoff, M., & Hoffken, O. (2016). Repetitive transcranial direct current stimulation induced excitability changes of primary visual cortex and visual learning effects: A pilot study. *Frontiers in Behavioral Neuroscience*, *10*(116), 116, <https://doi.org/10.3389/fnbeh.2016.00116>.
- Shalev, N., De Wandell, L., Dockree, P., Demeyere, N., & Chechlacz, M. (2018). Beyond time and space: The effect of a lateralized sustained attention task and brain stimulation on spatial and selective attention. *Cortex*, *107*, 131–147, <https://doi.org/10.1016/j.cortex.2017.09.009>.
- Terney, D., Chaieb, L., Moliadze, V., Antal, A., & Paulus, W. (2008). Increasing human brain excitability by transcranial high-frequency random noise stimulation. *Journal of Neuroscience*, *28*(52), 14147–14155, <https://doi.org/10.1523/JNEUROSCI.4248-08.2008>.
- Tsodyks, M., & Gilbert, C. (2004). Neural networks and perceptual learning. *Nature*, *431*(7010), 775–781, <https://doi.org/10.1038/nature03013>.
- Tyler, S. C., Conto, F., & Battelli, L. (2018). Rapid improvement on a temporal attention task within a single session of high-frequency transcranial random noise stimulation. *Journal of Cognitive Neuroscience*, *30*(5), 656–666, https://doi.org/10.1162/jocn_a_01235.
- Van Meel, C., Daniels, N., de Beeck, H. O., & Baeck, A. (2016). Effect of tDCS on task relevant and irrelevant perceptual learning of complex objects. *Journal of Vision*, *16*(6):, 1–15, <https://doi.org/10.1167/16.6.13>.
- Watson, A. B., & Pelli, D. G. (1983). QUEST: A Bayesian adaptive psychometric method. *Perception & Psychophysics*, *33*(2), 113–120, <https://doi.org/10.3758/bf03202828>.
- Wu, D., Wang, Y., Liu, N., Wang, P., Sun, K., & Xiao, W. (2022). High-definition transcranial direct current stimulation of the left middle temporal complex does not affect visual motion perception learning. *Frontiers in Neuroscience*, *16*, 988590, <https://doi.org/10.3389/fnins.2022.988590>.
- Wu, D., Wang, Y., Liu, N., Wang, P., Sun, K., & Zhang, P. (2023). Posttraining anodal tDCS

- improves early consolidation of visual perceptual learning. *Clinical Neurophysiology*, 146, 89–96, <https://doi.org/10.1016/j.clinph.2022.11.016>.
- Wu, D., Zhang, P., Wang, Y., Liu, N., Sun, K., Wang, P., . . . Xiao, W. (2023). Anodal online transcranial direct current stimulation facilitates visual motion perceptual learning. *European Journal of Neuroscience*, 57(3), 479–489, <https://doi.org/10.1111/ejn.15895>.
- Wu, S., Xin, C., Zhu, W., Wan, Y., Sang, Q., Du, R., . . . Wang, N. L. (2023). Transformation of cognition from mechanical to biological aqueous outflow pump may be a breakthrough in solving the problem of intraocular pressure regulation in glaucoma. *Science China Life Sciences*, 66(7), 1693–1696, <https://doi.org/10.1007/s11427-023-2342-8>.
- Yang, J., Yan, F. F., Chen, L., Xi, J., Fan, S., Zhang, P., . . . Huang, C. B. (2020). General learning ability in perceptual learning. *Proceedings of the National Academy of Sciences, USA*, 117(32), 19092–19100, <https://doi.org/10.1073/pnas.2002903117>.
- Yang, X. Y., He, Q., & Fang, F. (2022). Transcranial direct current stimulation over the visual cortex facilitates awake consolidation of visual perceptual learning. *Brain Stimulation*, 15(2), 380–382, <https://doi.org/10.1016/j.brs.2022.01.019>.