

Position shifts of fMRI-based population receptive fields in human visual cortex induced by Ponzo illusion

Dongjun He, Ce Mo, Yizhou Wang & Fang Fang

Experimental Brain Research


ISSN 0014-4819

Volume 233

Number 12

Exp Brain Res (2015) 233:3535–3541

DOI 10.1007/s00221-015-4425-3



Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Position shifts of fMRI-based population receptive fields in human visual cortex induced by Ponzo illusion


Dongjun He^{1,2,3,4} · Ce Mo³ · Yizhou Wang⁵ · Fang Fang^{1,2,3,4,48.27469802}

Received: 5 June 2015 / Accepted: 18 August 2015 / Published online: 28 August 2015
Springer-Verlag Berlin Heidelberg 2015

pRF position shift in human visual areas, especially in V1, lending further support for the receptive field position shift explanation for the Ponzo illusion.

Keywords Ponzo illusion · Population receptive field · Functional brain imaging

Fig. 1 A model of perceived size representation in early visual cortex. The perceived size is determined by the spatial distribution of activated neurons. The ring stimulus presented with no context (*left*) activates neurons through whose RFs it passes. The ring that appears to be close is perceived as small, consistent with outward RF shifts and a small cortical representation (*middle*). Conversely, the ring that appears to be far is perceived as large, consistent with inward RF shifts and a large cortical representation (*right*)



findings suggest that the Ponzo illusion is mediated by the topographic dynamics of V1 neural activity modulated by feedback signal from higher cortical areas.

However, it is still unclear what mechanisms drive the shifts of the spatial distribution of V1 activity during the Ponzo illusion. Neurons in V1 are characterized by their precise retinotopic organization. Their receptive field positions could serve to encode the spatial properties (e.g., location and size) of visual inputs. To represent the perceptually larger ring, neurons with more peripheral receptive fields need to be recruited to encode the perceived size of the ring. Hence, a possible neuronal mechanism in V1 for generating the Ponzo illusion is the position shift of V1 neuronal receptive fields initiated by feedback signal because visual depth information in a 3D scene is highly complex and extracting the information is beyond the local processing capacity of V1 neurons with limited receptive field sizes. In this case, the far ring elevates the firing rate of peripheral neurons as a result of an inward shift of the receptive fields of the neurons toward the fovea, leading to more peripheral V1 activity (Fig. 1). However, it is traditionally believed that the receptive field positions of V1 neurons are mainly determined by the pattern of feed forward connections from the retina to the cortex independent of current task demands and attention states (Alonso et al. 2001; Lund et al. 2003; Reid and Alonso 1995). Although a recent study reported neuronal receptive field position shifts in macaque primary visual cortex during the Ponzo illusion (Ni et al. 2014), so far there is no such evidence in human subjects because of the rare opportunity of intracranial recording in human early visual cortex (Mukamel and Fried 2012).

One approach to address this question is the non-invasive fMRI-based population receptive field (pRF) mapping technique (Dumoulin and Wandell 2008). This technique is based on the assumption that the joint receptive field of the neuronal population within a single voxel can be characterized by a two-dimensional Gaussian function with three parameters, x_0 , y_0 , and σ , where x_0 and y_0 determine the position (i.e., center) of the joint receptive field in the visual field and σ (i.e., dispersion of the Gaussian) determines the size of the receptive field. By fitting the predicted signal based on this model to the BOLD signal time course, the pRF position and size parameters can be estimated for individual voxels, thus providing a characterization of the receptive field properties of neuronal populations across the visual cortex. This method has been shown to reconstruct the cortical visual field map more accurately than conventional phase-encoded retinotopic mapping methods and produce pRF size estimates that agree well with electrophysiological measurements in monkey and human subjects (Kok and de Lange 2014).

In the current study, we used the fMRI-based pRF mapping technique in combination of psychophysics to investigate whether the Ponzo illusion is mediated by systematic shifts of pRF positions in human early visual cortex. The objective of the psychophysical experiment was to measure the magnitude of the Ponzo illusion, and the pRF mapping experiment was designed to measure the pRF properties in human early visual areas during the Ponzo illusion. We used similar checkered ring stimuli and 3D visual scene as those in Fang et al. (2008) (Fig. 2a). We hypothesized that, relative to the close ring, the depth cues in the 3D scene would cause systematic inward shifts of the pRF positions




Fig. 2 Stimuli and experimental procedure. **a** Stimuli. One stimulus was a rendered 3D scene of a hallway and walls with two physically identical rings: one ring was at a close apparent depth and the other was at a far apparent depth. In the other stimulus, the two rings were

presented against a gray background. **b** fMRI experimental procedure. Expanding and contracting rings were presented at either the close or far apparent depth for estimating pRF parameters in V1–V3

of the voxels responding to the far ring area, leading to a larger perceived size of the far ring.

Materials and methods

Subjects

A total of nine human subjects (6 male, 18–28 years old) were paid to take part in the study. All of them participated in both the psychophysical and fMRI experiments. All subjects were naive to the purpose of the study. They were right-handed, reported normal or corrected-to-normal vision, and had no known neurological or visual disorders. They gave written informed consent, and our procedures were approved by the Human Subject Review Committee at Peking University.

Psychophysical experiment

The psychophysical experiment was performed in a separate session outside the magnet. Visual stimuli were presented on an IIYAMA monitor (model HM204DT, size 22 inches) with a spatial resolution of 1024×768 and a refresh rate of 85 Hz. Subjects viewed the stimuli from a distance of 73 cm with their head stabilized on a chin rest. We used a method of constant stimuli to measure the magnitude of the Ponzo illusion. We presented two rings in a rendered three-dimensional (3D) scene of a hallway and walls: one ring was at a close (“front”) apparent depth and the other one was at a far (“back”) apparent depth

(Fig. 2a). In a trial, subjects were given 2 s of free viewing time before the rings disappeared and then were required to indicate which ring looked larger. The size of the front ring was fixed. Its inner and outer radii were 1.96° and 2.52° , respectively. The size of the back ring varied from trial to trial. Its inner and outer radii were randomly selected from five pairs of values: $1.40^\circ/1.96^\circ$, $1.68^\circ/2.24^\circ$, $1.96^\circ/2.52^\circ$, $2.24^\circ/2.80^\circ$, and $2.52^\circ/3.08^\circ$. Note that all the front and back rings had a fixed width (0.56°). Subjects completed 200 trials in total, with 40 trials for each back ring size. In addition to the 3D scene condition, we also presented the rings on a gray background with no contextual cue and measured the perceived size difference between the rings, serving as a baseline for comparison (Fig. 2a). For each of the two background conditions, we plotted the percentage of trials in which subjects indicated that the back ring was perceived to be larger than the front ring as a function of the physical size of the back ring. To quantitatively measure the illusion size, we fit the psychometric values at the five physical sizes with a cumulative normal function. We interpolated the data to find the point of subjective equality (PSE) at which the front and back rings appeared to have the same size.

fMRI experiments

Retinotopic visual areas (V1, V2, and V3) were defined by a standard phase-encoded method developed by Sereno et al. (1995) and Engel et al. (1997), in which subjects viewed a rotating wedge and an expanding ring that created traveling waves of neural activity in visual cortex.

An independent block-design run was performed to identify the voxels or ROIs in the retinotopic areas responding to the front and back rings when subjects fixated at the center of the rings. The run contained eight stimulus blocks of 12 s, interleaved with eight blank blocks of 12 s. The stimulus was a full-contrast flickering checkered ring (inner radius 1.42° , outer radius 3.10°).

To estimate pRF parameters, we measured hemodynamic response function (HRF) for each subject in a separate run. This run contained 12 trials. In each trial, a full-contrast flickering checkered disc with a radius of 10.94° was presented for 2 s, followed by a 30-s blank interval. The HRF was measured by fitting the convolution of a 6-parameter double-gamma function with a 2-s boxcar function to the BOLD response elicited by the disc. In both the localizer run and the HRF estimation run, subjects performed a color discrimination task at fixation point to maintain fixation and control attention.

The pRF mapping experiment consisted of 8 functional runs of 192 s (Fig. 2b). In half of the runs, a series of flickering checkered rings with 16 different sizes (inner/outer radius of the smallest ring $0^\circ/0.56^\circ$, inner/outer radius of the largest ring $4.2^\circ/4.76^\circ$, step size 0.28°) were presented at the front ring position, while in the other half, the ring




Fig. 3 Psychophysical and fMRI results. **a** The magnitudes of the Ponzo illusion in the 3D scene condition and in the *gray* background condition. The *dashed line* indicates the outer radius of the front ring. The PSE was the outer radius of the back ring when the front and back rings were perceived to be equally large. **b** pRF positions in V1–

V3 mapped by the front and back rings. *Asterisks* indicate a statistically significant difference between the front and back ring conditions (** $p < 0.01$; *** $p < 0.001$). *Error bars* represent 1 SEM across subjects




Fig. 4 Correlations between the magnitude of the Ponzo illusion and the pRF position shift in V1–V3 across individual subjects

of 1.81° and 2.37° (5.95 % of the outer radius of the front ring), the two rings were perceived to be equally large. The magnitude of the Ponzo illusion was quantified as the size difference between the two rings when the PSE was identified (mean \pm sem $0.15^\circ \pm 0.02^\circ$). The illusion size was robust and significant [$t(8) = 8.68$, $p < 0.001$] (Fig. 3a). It is noteworthy that, when the front and back rings were presented on a gray background, the illusion completely vanished (0.07% of the outer radius of the front ring), suggesting that the Ponzo illusion observed in the 3D scene condition was not simply due to the locations of the rings (Fig. 3a).

fMRI results

We estimated the pRF positions of the voxels corresponding to the front and back rings, respectively. The estimated

positions were converted to eccentricities and then averaged across voxels in V1, V2, or V3 (Fig. 3b). In V1, the mean pRF eccentricities associated with the front and back rings were 2.34° and 2.18° , respectively. This finding demonstrated that the far depth information, relative to the near depth information, caused an inward pRF shift toward the fovea [$t(8) = 4.09$, $p < 0.005$]. A similar pattern was found in the other two visual areas. In V2, the mean pRF eccentricities associated with the front and back rings were 2.35° and 2.20° , respectively [$t(8) = 4.33$, $p < 0.005$]. In V3, the mean pRF eccentricities associated with the front and back rings were 2.38° and 2.24° , respectively [$t(8) = 5.15$, $p < 0.001$].

To further examine the behavioral relevance of the observed pRF position shifts, we calculated the Pearson's correlation coefficient between the magnitude of the Ponzo illusion and the pRF position shifts in V1–V3 (Fig. 4). We

found a significant positive correlation in V1 ($r = 0.92$, $p < 0.001$), but not in V2 ($r = 0.569$, $p = 0.11$, 32 % variance explained) and V3 ($r = 0.64$, $p = 0.063$, 41 % variance explained). The correlation coefficient in V1 was (marginally) significantly larger than those in V2 ($p < 0.05$) and V3 ($p = 0.07$), suggesting that V1 was more closely associated with the Ponzo illusion than V2 and V3.

Discussion

Our study found that, relative to the close apparent depth in the 3D scene, the far apparent depth caused the pRF positions of voxels in V1–V3 to shift toward the fovea, consistent with the perceived size of the front and back rings. Moreover, the pRF position shift in V1 significantly correlated with the magnitude of the Ponzo illusion across individual subjects. It should be noted that, the observed effects are unlikely due to difference in spatial attention or eye movement, because subjects performed the same task with the front and back rings.

Fang et al. (2008) revealed that the Ponzo illusion could alter the spatial pattern of V1 neural activity and the perceived larger ring activated more peripheral areas in V1, compared with the perceived smaller ring. To represent the perceptually larger ring, neurons with more peripheral receptive fields need to be recruited to encode the perceived size of the back ring. The observed inward shift of the pRFs toward the fovea entails more peripheral neurons to represent the perceived size of the back ring, leading to more peripheral V1 activity. Thus, our finding here is in line with Fang et al. (2008). Using extracellular recordings, Ni et al. (2014) also found a similar receptive field shift of V1 neurons. Our study here examined not only V1, but also extrastriate areas V2 and V3, thus providing a more comprehensive characterization of neural changes during the Ponzo illusion. Notably, our observation of the systematic receptive field shifts in V1–V3 is in line with the neurological zoom system account for perceptual size constancy proposed in an early study (Marg and Adams 1970). Using the technique of implanted micro-electrode recording, the researchers found that the neuronal receptive fields in human early visual cortex were “magnified” in response to an incoming object such that the perceived object size was not altered by the increased retinal image. Together, these studies make headways toward unraveling the neural mechanisms of the Ponzo illusion.

Consistent with our results, recent studies also demonstrated that the topography of V1 neural activity closely mirrored the perceived object size that was significantly affected by spatiotemporal contexts (e.g., physical distance or the size of visual adaptor), even when the retinal input remains unchanged (Pooresmaeili et al. 2013; Sperandio

et al. 2012). In the study conducted by Sperandio et al. (2012), they presented subjects with a constant light stimulus and manipulated the perceived size of the stimulus afterimage by varying the viewing distance of a back screen holding the afterimage. They found that the spatial distribution of V1 activity expanded to the more eccentric representation of the visual field as the perceived size of the afterimage increased with the viewing distance, suggesting neuronal populations in V1 encode the size information of the perceived rather than the physical visual input. In another study performed by Pooresmaeili et al. (2013), using a visual adaptation paradigm, they found that the spatial extent of V1 activation closely resembled the perceived size of the testing stimulus that was altered by the size of adapting stimulus. Moreover, other recent studies also demonstrated that neural activity in V1 could encode other perceived stimulus properties (e.g., brightness and shape; Boyaci et al. 2007; Michel et al. 2013). Together, these findings provide strong evidence of the critical role of V1 in encoding perceived (rather than physical) visual information, which is usually believed to take place in high visual areas.

Across individual subjects, the pRF shift in V1 exhibited a stronger coupling with the magnitude of the Ponzo illusion than those in V2 and V3. One possible explanation is that the integration of size and distance information for size–distance scaling is dependent on top-down modulation that is better captured by the BOLD responses in striate than in extrastriate areas (Sperandio and Chouinard 2015). This speculation is in accordance with a recent anatomical finding (Schwarzkopf et al. 2011). Schwarzkopf et al. (2011) found that the magnitude of the Ponzo illusion was negatively correlated with the surface area of V1 but not with that of V2 and V3. Together, these findings reveal the functional and anatomical bases underlying the interindividual difference in susceptibility to the Ponzo illusion. How does our functional finding relate to Schwarzkopf et al.’s anatomical finding? Horizontal connections in V1 provide an effective means for individual neurons to integrate and assess visual information over a large portion of the visual field beyond their own receptive fields (Gilbert and Wiesel 1989), which might be critical for the receptive field shift of V1 neurons observed here. As the area of V1 surface increases, the horizontal connections become less effective as the information has to travel longer distances for interneuron communication, leading to smaller receptive field shifts. Hence, we speculate that horizontal connections in V1 might serve as an important predictor of individual susceptibility to the Ponzo illusion.

In sum, we show that the neural coding of perceived object sizes during the Ponzo illusion is implemented in the human brain via systematic shifts of pRF positions throughout early visual areas, which provides the non-invasive

model-based fMRI evidence in human subjects echoing previous invasive electrophysiological findings in monkey subjects.

Acknowledgments This work was supported by the Ministry of Science and Technology of China (2015CB351800) and the National Natural Science Foundation of China (31230029 and 31421003). We thank the staff at the Center for MRI Research at Peking University for technical support.

References

- Alonso JM, Usrey WM, Reid RC (2001) Rules of connectivity between geniculate cells and simple cells in cat primary visual cortex. *J Neurosci* 21(11):4002–4015
- Boyaci H, Fang F, Murray SO, Kersten D (2007) Responses to lightness variations in early human visual cortex. *Curr Biol* 17(11):989–993
- Dumoulin SO, Wandell BA (2008) Population receptive field estimates in human visual cortex. *Neuroimage* 39(2):647–660
- Engel SA, Glover GH, Wandell BA (1997) Retinotopic organization in human visual cortex and the spatial precision of functional MRI. *Cereb Cortex* 7(2):181–192
- Fang F, Boyaci H, Kersten D, Murray SO (2008) Attention-dependent representation of a size illusion in human V1. *Curr Biol* 18(21):1707–1712
- Gilbert CD, Wiesel TN (1989) Columnar specificity of intrinsic horizontal and corticocortical connections in cat visual cortex. *J Neurosci* 9(7):2432–2442
- Kok P, de Lange FP (2014) Shape perception simultaneously up- and downregulates neural activity in the primary visual cortex. *Curr Biol* 24(13):1531–1535
- Leibowitz H, Brislin R, Perlmutter L, Hennessy R (1969) Ponzo perspective illusion as a manifestation of space perception. *Science* 166(3909):1174–1176
- Lund JS, Angelucci A, Bressloff PC (2003) Anatomical substrates for functional columns in macaque monkey primary visual cortex. *Cereb Cortex* 13(1):15–24
- Marg E, Adams JE (1970) Evidence for a neurological zoom system in vision from angular changes in some receptive fields of single neurons with changes in fixation distance in the human visual cortex. *Experientia* 26(3):270–271
- Michel MM, Chen Y, Geisler WS, Seidemann E (2013) An illusion predicted by V1 population activity implicates cortical topography in shape perception. *Nat Neurosci* 16(10):1477–1483
- Mukamel R, Fried I (2012) Human intracranial recordings and cognitive neuroscience. *Ann Rev Psychol* 63:511–537
- Murray SO, Boyaci H, Kersten D (2006) The representation of perceived angular size in human primary visual cortex. *Nat Neurosci* 9(3):429–434
- Ni AM, Murray SO, Horwitz GD (2014) Object-centered shifts of receptive field positions in monkey primary visual cortex. *Curr Biol* 24(14):1653–1658
- Pooresmaeili A, Arrighi R, Biagi L, Morrone MC (2013) Blood oxygen level-dependent activation of the primary visual cortex predicts size adaptation illusion. *J Neurosci* 33(40):15999–16008
- Reid RC, Alonso JM (1995) Specificity of monosynaptic connections from thalamus to visual cortex. *Nature* 378(6554):281–283
- Schwarzkopf DS, Song C, Rees G (2011) The surface area of human V1 predicts the subjective experience of object size. *Nat Neurosci* 14(1):28–30
- Sereno MI, Dale AM, Reppas JB, Kwong KK, Belliveau JW, Brady TJ et al (1995) Borders of multiple visual areas in humans revealed by functional magnetic resonance imaging. *Science* 268(5212):889–893
- Sperandio I, Chouinard PA (2015) The mechanisms of size constancy. *Multisens Res* 28(3–4):253–283
- Sperandio I, Chouinard PA, Goodale MA (2012) Retinotopic activity in V1 reflects the perceived and not the retinal size of an afterimage. *Nat Neurosci* 15(4):540–542